Self-Monitoring Blood Glucose (SMBG) Frequency & Pattern Tool

http://guidelines.diabetes.ca/self-management/smbg-tool

Pattern Recommendation: Diagnosis of type 2 diabetes in the past 6 months or not meeting glycemic targets

Check \geq 1 time per day, on average, and at different times of day to learn the effects of various meals, exercise and/or medications on blood glucose.

Please pick one of the following SMBG patterns below

SMBG basic pattern

	breakfast		lunch		supper		- bedtime	night
	before	after	before	after	before	after	bedume	riigric
Sunday								
Monday								
Tuesday								
Wednesday								
Thursday								
Friday								
Saturday								

SMBG pattern to learn the effects of various meals

	breakfast		lunch		supper		bedtime	night
	before	after	before	after	before	after	bedume	riigric
Sunday								
Monday								
Tuesday								
Wednesday								
Thursday								
Friday								
Saturday								



guidelines.diabetes.ca | 1-800 BANTING (226-8464) | diabetes.ca

Self-Monitoring Blood Glucose (SMBG) Frequency & Pattern Tool

http://guidelines.diabetes.ca/self-management/smbg-tool

Pattern Recommendation (continued): Diagnosis of type 2 diabetes in the past 6 months or not meeting glycemic targets

SMBG pattern to learn the effects of exercise

• Sometimes benefit is seen 4 – 8 hours later, so alternate post-exercise testing times.

	break	ĸfast	lunch		supper		bedtime	night	
	before	after	before	after	before	after	bedtime	Tilgitt	
Sunday				Exercise	TEST				
Monday									
Tuesday									
Wednesday									
Thursday						Exercise	TEST		
Friday									
Saturday									

SMBG to get a snapshot of blood glucose levels throughout the day, on one day of the week, which is a typical day.

	breakfast		lunch		supper		- bedtime	night
	before	after	before	after	before	after	beduine	Iligilic
Sunday								
Monday								
Tuesday								
Wednesday								
Thursday								
Friday								
Saturday								



guidelines.diabetes.ca | 1-800 BANTING (226-8464) | diabetes.ca